

Appropriate laboratory testing for methamphetamine exposure

Reference list compiled by:

Grant D. Beardsley, M.S., MT (ASCP)
Clinical Toxicologist, Oregon Medical Laboratories
Manager, Toxicology
541-686-8137
gbeardsley@omlabs.com

Suggested references:

Stout PR, Horn CK, Klette KL. *Rapid simultaneous determination of amphetamine, methamphetamine, 3,4-methylenedioxyamphetamine, 3,4-methylenedioxy-methamphetamine, and 3,4-methylenedioxyethylamphetamine in urine by solid-phase extraction and GC-MS: a method optimized for high-volume laboratories.* J Anal Toxicol. 2002 Jul-Aug;26(5):253-61.

Valentine JL, Kearns GL, Sparks C, Letzig LG, Valentine CR, Shappell SA, Neri DF, DeJohn CA. *GC-MS determination of amphetamine and methamphetamine in human urine for 12 hours following oral administration of dextro-methamphetamine: lack of evidence supporting the established forensic guidelines for methamphetamine confirmation.* J Anal Toxicol. 1995 Nov-Dec;19(7):581-90. Department of Pediatrics, University of Arkansas for Medical Sciences and Toxicology, Arkansas Children's Hospital, Little Rock 72202-3591, USA.

Gospe SM Jr. *Transient cortical blindness in an infant exposed to methamphetamine.* Ann Emerg Med. 1995 Sep;26(3):380-2. Department of Neurology, University of California, Davis, Medical Center, Sacramento, USA.

Arria AM, Derauf C, Lagasse LL, Grant P, Shah R, Smith L, Haning W, Huestis M, Strauss A, Grotta SD, Liu J, Lester B. *Methamphetamine and other substance use during pregnancy: preliminary estimates from the Infant Development, Environment, and Lifestyle (IDEAL) study.* Matern Child Health J. 2006 May;10(3):293-302. Epub 2006 Jan 5. Center for Substance Abuse Research, University of Maryland, College Park, Maryland.

Montgomery D, Plate C, Alder SC, Jones M, Jones J, Christensen RD. *Testing for fetal exposure to illicit drugs using umbilical cord tissue vs meconium.* J Perinatol. 2006 Jan 1;26(1):11-4. Department of Women and Newborns, Intermountain Health Care and McKay Dee Hospital, Ogden, UT 84403, and Department of Family and Preventive Medicine, University of Utah, Salt Lake City, UT, USA.

Casanova OQ, Lombarbadero N, Behnke M, Eyler FD, Conlon M, Bertholf RL. *Detection of cocaine exposure in the neonate. Analysis of urine, meconium and amniotic fluid from mothers and infants exposed to cocaine.* Arch Pathol Lab Med 1994; 118:988-993. Lawnwood Regional Medical Center, Ft Pierce, Florida; the departments of Pathology and Lab Medicine, Pediatrics and Biostatistics, University of Florida College of Medicine, Gainesville, Florida.

Kwong TC, Ryan RM. *Detection of intrauterine illicit drug exposure by newborn drug testing.* Clin Chem 1997; 43(1):235-242. Departments of Pathology and Laboratory Medicine and Pediatrics, School of Medicine, University of Rochester, NY.

Moriya F, Chan KM, Noguchi TT, Wu PK. *Testing for drugs of abuse in meconium of newborn infants.* J Anal Toxicol. 1994 Jan-Feb;18(1):41-45. Department of Laboratories and Pathology and Division of Neonatal-Perinatal Medicine, Department of Pediatrics, University of Southern California Medical Center, Los Angeles, California.